



UPWARD SMILES, INC.
www.upwardsmiles.com

Extraction Post-Op Instructions

One or more teeth have been 'wiggled' out today, and it is important for the patient to follow these instructions to allow the mouth to heal quickly.

Today: Continue biting on the gauze for 30-60 minutes following your procedure. Do NOT spit, rinse, suck (using a straw), smoke, drink carbonated or alcoholic beverages for at least 24 hours. Passively empty your mouth when needed.

- The goal is to keep blood clot in the socket to prevent bleeding and a painful complication known as DRY SOCKET.

Day 2: Patient will most likely experience one or more of the following symptoms:

- Minor pain, swelling, bruising, limitation in opening or closing of the mouth, pain in jaw or ear, sore throat. It is important for the patient to listen to their body and take it easy.

No heavy lifting, spitting, sucking, alcohol or smoking.

Oral hygiene: Brush gently. A gentle lukewarm salt water rinse can be used to freshen your mouth. But again, do not spit; just let the fluid passively empty. The corners of your mouth may become cracked and dry – moisturize frequently.

Discomfort: Over-the-counter- pain medication (Ibuprofen, Aleve, and Tylenol) may be taken as directed on the bottle to relieve discomfort.

Swelling: Apply cold compresses to affected area at 10 minute intervals for the first 24 hours to minimize swelling. Any swelling that occurs usually begins to diminish within 72 hours; call the office if there is no change.

Bleeding: Some oozing of blood is normal for the first 12-24 hours. Put a hand towel on your pillow as some drooling can occur when you are numb. If you experience excessive bleeding, apply firm pressure with 1-2 folded gauze pads or damp tea bag on the affected area for 30-60 minutes and keep your head elevated. Call the office if the bleeding does not subside.

Diet: Drink plenty of fluids, avoid carbonated and alcoholic beverages. A soft diet will be easiest: yogurt, milkshakes (no straw), soups, fish, pasta. Nothing too spicy is recommended. It is normal to have a little tenderness while chewing and difficulty opening wide.

Prescriptions: If you have been prescribed antibiotics, take them as directed **until they are all gone**, even if symptoms dissipate. If pain medication or a mouth rinse is prescribed take as directed.

Should you have further questions, or if the patient experiences persistent pain, bleeding, or swelling for more than two days, please contact our office at 855-944-5437.